

UC Santa Cruz

UC Santa Cruz Previously Published Works

Title

Genome Sequence of the Photoarsenotrophic Bacterium Ectothiorhodospira sp. Strain BSL-9, Isolated from a Hypersaline Alkaline Arsenic-Rich Extreme Environment.

Permalink

<https://escholarship.org/uc/item/7gt31097>

Journal

Genome announcements, 4(5)

ISSN

2169-8287

Authors

Hernandez-Maldonado, Jaime
Stoneburner, Brendon
Boren, Alison
et al.

Publication Date

2016-10-01

DOI

10.1128/genomea.01139-16

Peer reviewed

Genome Sequence of the Photoarsenotrophic Bacterium *Ectothiorhodospira* sp. Strain BSL-9, Isolated from a Hypersaline Alkaline Arsenic-Rich Extreme Environment

Jaime Hernandez-Maldonado,^a Brendon Stoneburner,^a Alison Boren,^a Laurence Miller,^b Michael Rosen,^c Ronald S. Oremland,^b Chad W. Saltikov^a

Department of Microbiology and Environmental Toxicology, University of California Santa Cruz, Santa Cruz, California, USA^a; U.S. Geological Survey, Menlo Park, California, USA^b; U.S. Geological Survey, Carson City, Nevada, USA^c

The full genome sequence of *Ectothiorhodospira* sp. strain BSL-9 is reported here. This purple sulfur bacterium encodes an *arxA*-type arsenite oxidase within the *arxB2AB1CD* gene island and is capable of carrying out “photoarsenotrophy” anoxygenic photosynthetic arsenite oxidation. Its genome is composed of 3.5 Mb and has approximately 63% G+C content.

Received 23 August 2016 Accepted 25 August 2016 Published 13 October 2016

Citation Hernandez-Maldonado J, Stoneburner B, Boren A, Miller L, Rosen M, Oremland RS, Saltikov CW. 2016. Genome sequence of the photoarsenotrophic bacterium *Ectothiorhodospira* sp. strain BSL-9, isolated from a hypersaline alkaline arsenic-rich extreme environment. *Genome Announc* 4(5):e01139-16. doi:10.1128/genomeA.01139-16.

Copyright © 2016 Hernandez-Maldonado et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Chad W. Saltikov, saltikov@ucsc.edu.

Arsenic-rich soda lakes are ideal environments for culturing microorganisms with unique metabolic capabilities for coupling cellular energy production to arsenic oxidation and reduction (1–6). Here, we report the assembled genome of an anoxygenic photosynthetic arsenite-oxidizing (“photoarsenotrophic”) bacterium, *Ectothiorhodospira* sp. strain BSL-9. This microbe was isolated from Big Soda Lake, an arsenic-rich (~25 μM), hypersaline (26 to 88 g/liter total dissolved solids), alkaline (pH 9.7) lake located in Nevada (39°31'N 118°52'W) (7–9). Moreover, this crater lake has a well-defined seasonal bloom of purple sulfur bacteria (*Chromatium* and *Ectothiorhodospira* species) (10) that are proposed to contribute to the arsenic geochemical cycle.

Assessment of the BSL-9 genome revealed an arsenic gene island, *arxB2AB1CD* (11), which is predicted to encode the arsenite oxidase gene *arxA*. Moreover, *arxB2AB1CD* encodes a [4Fe-4S]-containing protein (*arxB2*), a second [4Fe-4S]-containing protein (*arxB1*), a membrane protein (*arxC*), and a TorD-like protein involved in molybdenum enzyme biogenesis (*arxD*). In the chemolithotrophic bacterium *Alkalilimnicola ehrlichii* sp. strain MLHE-1, *arxA* is required for anaerobic arsenite oxidation coupled to nitrate (12). The BSL-9 genome lacks the AioA-type arsenite oxidase. A BSL-9 *arxA* mutant strain shows that *arxA* is the sole arsenite oxidase for photoarsenotrophy (13).

Ectothiorhodospiraceae are common anoxygenic phototrophs with versatile abilities to metabolize inorganic and organic electron donors (14–16), which enables them to occupy distinct euphotic hypersaline alkaline environments. In addition to arsenite, BSL-9 can grow photoautotrophically with sulfide or thiosulfate. This is consistent with the presence of *sox* and *dsr* genes, which are involved in sulfur oxidation (17, 18). Moreover, BSL-9 can also grow as a photoheterotroph with various organic acids (e.g., acetate, malate, propionate, lactate, fumarate, succinate, and pyruvate). BSL-9 is sensitive to chloramphenicol, resistant to kanamycin, carbenicillin, gentamicin, and tetracycline, and grows optimally at 35°C at pH 8, 2% NaCl; these growth patterns are consistent with other

Ectothiorhodospira species (14–16). Although BSL-9 is an anaerobe, the presence of cytochrome *c* oxidase genes (e.g. *cbb₃*) found in BSL-9 may explain its tolerance to atmospheric oxygen. For example, cytochrome *c* oxidases (*cbb₃*) are known for having high oxygen affinity, and cytochrome *c* peroxidases protect cells from reactive oxygen species. The BSL-9 genome also encodes photosynthetic complex genes, such as bacteriochlorophyll *a* synthase, the light-harvesting complex *pucAB*, and two copies of the carbon fixation-related gene *rbcL* (type III RuBisCO). Having the full genome sequence of BSL-9 opens numerous possibilities for studying the metabolic abilities, physiology, and the ecological environmental impact of photoarsenotrophy to the arsenic biogeochemical cycle.

The genome was done at the UC Davis Genome Sequencing Center using PacBio technology. The Hierarchical Genome Assembly Process (HGAP_v2) assembly pipeline (19) was used with ~300× sequence coverage. For annotation, the NCBI Public Genome Annotation Pipeline service was used. The resulting assembly was 3.5 Mb, with 63% G+C content.

Accession number(s). The genome sequence of *Ectothiorhodospira* sp. strain BSL-9 was deposited in the GenBank database under the accession no. [CP011994](https://www.ncbi.nlm.nih.gov/nuclot/CP011994), NCBI BioProject accession no. PRJNA232800, and BioSample accession no. SAMN03795182.

ACKNOWLEDGMENT

We thank the Sierra Nevada Aquatic Research Laboratory for facilitating the fieldwork.

FUNDING INFORMATION

This work, including the efforts of Jaime Hernandez-Maldonado, was funded by NIH-NHGRI (1R25HG006836-01A1 and 5R25GM058903-14). This work, including the efforts of Chad W. Saltikov, was funded by National Science Foundation (NSF) (EAR-1349366) and the Water Mission Area of the U.S. Geological Survey.

REFERENCES

- Oremland RS, Kulp TR, Blum JS, Hoeft SE, Baesman S, Miller LG, Stolz JF. 2005. A microbial arsenic cycle in a salt-saturated, extreme environment. *Science* 308:1305–1308. <http://dx.doi.org/10.1126/science.1110832>.
- Lloyd JR, Oremland RS. 2006. Microbial transformations of arsenic in the environment: from soda lakes to aquifers. *Elements* 2:85–90. <http://dx.doi.org/10.2113/gselements.2.2.85>.
- Hoeft SE, Blum JS, Stolz JF, Tabita FR, Witte B, King GM, Santini JM, Oremland RS. 2007. *Alkalilimnicola ehrlichii* sp. nov., a novel, arsenite-oxidizing haloalkaliphilic gammaproteobacterium capable of chemoautotrophic or heterotrophic growth with nitrate or oxygen as the electron acceptor. *Int J Syst Evol Microbiol* 57:504–512. <http://dx.doi.org/10.1099/ijs.0.64576-0>.
- Kulp TR, Hoeft SE, Asao M, Madigan MT, Hollibaugh JT, Fisher JC, Stolz JF, Culbertson CW, Miller LG, Oremland RS. 2008. Arsenic(III) fuels anoxygenic photosynthesis in hot spring biofilms from Mono Lake, California. *Science* 321:967–970. <http://dx.doi.org/10.1126/science.1160799>.
- Stolz JF, Basu P, Oremland RS. 2010. Microbial arsenic metabolism: New twists on an old poison. *Microbe* 5:53–59. <http://dx.doi.org/10.1128/microbe.5.53.1>.
- Edwardson CF, Planer-Friedrich B, Hollibaugh JT. 2014. Transformation of monothioarsenate by haloalkaliphilic, anoxygenic photosynthetic purple sulfur bacteria. *FEMS Microbiol Ecol* 90:858–868. <http://dx.doi.org/10.1111/1574-6941.12440>.
- Cloern JE, Cole BE, Oremland RS. 1983. Seasonal changes in the chemistry and biology of a meromictic lake (Big Soda Lake, Nevada, USA). *Hydrobiologia* 105:195–206. <http://dx.doi.org/10.1007/BF00025188>.
- Cloern JE, Cole BE, Oremland RS. 1983. Autotrophic processes in meromictic Big Soda Lake, Nevada. *Limnol Oceanogr* 28:1049–1061. <http://dx.doi.org/10.4319/lo.1983.28.6.1049>.
- Priscu JC, Axler RP, Carlton RG, Reuter JE, Arneson PA, Goldman CR. 1982. Vertical profiles of primary productivity, biomass and physicochemical properties in meromictic Big Soda Lake, Nevada, USA. *Hydrobiologia* 96:113–120. <http://dx.doi.org/10.1007/BF02185426>.
- Zehr JP, Harvey RW, Oremland RS, Cloern JE, George LH, Lane JL. 1987. Big Soda Lake (Nevada). 1. Pelagic bacterial heterotrophy and biomass. *Limnol Oceanogr* 32:781–793. <http://dx.doi.org/10.4319/lo.1987.32.4.0781>.
- Zargar K, Conrad A, Bernick DL, Lowe TM, Stolz V, Hoeft S, Oremland RS, Stolz J, Saltikov CW. 2012. ArxA, a new clade of arsenite oxidase within the DMSO reductase family of molybdenum oxidoreductases. *Environ Microbiol* 14:1635–1645. <http://dx.doi.org/10.1111/j.1462-2920.2012.02722.x>.
- Zargar K, Hoeft S, Oremland R, Saltikov CW. 2010. Identification of a novel arsenite oxidase gene, *arxA*, in the haloalkaliphilic, arsenite-oxidizing bacterium *Alkalilimnicola ehrlichii* strain MLHE-1. *J Bacteriol* 192:3755–3762. <http://dx.doi.org/10.1128/JB.00244-10>.
- Hernandez-Maldonado J, Sanchez-Sedillo B, Stoneburner B, Boren A, Miller L, McCann S, Rosen M, Oremland RS, Saltikov CW. 2016. The genetic basis for anoxygenic photosynthetic arsenite oxidation. *Environ Microbiol* in press. <http://dx.doi.org/10.1111/1462-2920.13509>.
- Trüper HG. 1968. *Ectothiorhodospira mobilis* Pelsh, a photosynthetic sulfur bacterium depositing sulfur outside the cells. *J Bacteriol* 95:1910–1920.
- Imhoff JF, Tindall BJ, Grant WD, Truper HG. 1981. *Ectothiorhodospira vacuolata* sp. nov., a new phototrophic bacterium from soda lakes. *Arch Microbiol* 130:238–242. <http://dx.doi.org/10.1007/BF00459526>.
- Gorlenko VM, Bryantseva IA, Rabold S, Tourova TP, Rubtsova D, Smirnova E, Thiel V, Imhoff JF. 2009. *Ectothiorhodospira variabilis* sp. nov., an alkaliphilic and halophilic purple sulfur bacterium from soda lakes. *Int J Syst Evol Microbiol* 59:658–664. <http://dx.doi.org/10.1099/ijs.0.004648-0>.
- Friedrich CG, Bardischewsky F, Rother D, Quentmeier A, Fischer J. 2005. Prokaryotic sulfur oxidation. *Curr Opin Microbiol* 8:253–259. <http://dx.doi.org/10.1016/j.mib.2005.04.005>.
- Frigaard N-U, Dahl C. 2009. Sulfur metabolism in phototrophic sulfur bacteria. *Adv Microb Physiol* 54:103–200. [http://dx.doi.org/10.1016/S0065-2911\(08\)00002-7](http://dx.doi.org/10.1016/S0065-2911(08)00002-7).
- Chin C-S, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, Clum A, Copeland A, Huddleston J, Eichler EE, Turner SW, Korlach J. 2013. Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nat Methods* 10:563–569. <http://dx.doi.org/10.1038/nmeth.2474>.